

## I. Finding Roots of Quadratic Equations

- The Standard Form of a quadratic equation is:  $ax^2 + bx + c = 0$ .
- We can use the Quadratic Formula to solve equations in standard form:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

- Discriminant – The radical portion of this formula  $\sqrt{b^2 - 4ac}$ , determines the nature of the roots. This quantity under the radical sign  $b^2 - 4ac$ , is called the discriminant.

The diagram shows the quadratic formula  $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ . An arrow labeled "Radical" points to the square root symbol  $\sqrt{\phantom{x}}$ . Another arrow labeled "Discriminant" points to the expression  $b^2 - 4ac$  inside the square root.

- Three things may occur regarding the discriminant:
  - If  $b^2 - 4ac > 0$   
We can take the square root of this positive amount and there will be two different real answers (or roots) to the equation.
  - If  $b^2 - 4ac < 0$   
We cannot take the square root of a negative number, so there will be no real roots.
  - If  $b^2 - 4ac = 0$   
The amount under the radical is zero and since the square root of zero is zero, we will get only 1 distinct real root.

## II. Examples

a.  $x^2 - 6x + 9 = 0$   
 $a = 1$   $b = -6$   $c = 9$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{+6 \pm \sqrt{(-6)^2 - 4(1)(9)}}{2(1)}$$

$$x = \frac{6 \pm \sqrt{36 - 36}}{2}$$

$$x = \frac{6 \pm \sqrt{0}}{2} \leftarrow \boxed{\text{The Discriminant Equals Zero.}}$$

$$x = \frac{6 \pm 0}{2}$$

$$x = \frac{6}{2} = 3 \text{ (There is only 1 real root.)}$$

b.  $x^2 + 3x + 1 = 0$   
 $a = 1$   $b = 3$   $c = 1$

$$x = \frac{-3 \pm \sqrt{9 - 4(1)(1)}}{2(1)} \leftarrow \boxed{\text{The Discriminant is positive.}}$$

$$x = \frac{-3 \pm \sqrt{5}}{2}$$

$$x = \frac{-3 + \sqrt{5}}{2} \text{ and } x = \frac{-3 - \sqrt{5}}{2}$$

Since the discriminant is positive (it equals +5) there are two real roots.

c.  $x^2 + x + 3 = 0$   
 $a = 1$   $b = 1$   $c = 3$

$$x = \frac{-1 \pm \sqrt{1 - 4(1)(3)}}{2(1)}$$

The Discriminant is negative.

$$x = \frac{-1 \pm \sqrt{-11}}{2}$$

The discriminant is -11. Since we cannot take the square root of a negative number we have no real roots.

### III. Practice Problems

By examining the discriminant  $= b^2 - 4ac$ , determine how many real roots, if any, the following quadratic equations have.

1.  $x^2 - 4x + 4 = 0$

2.  $x^2 + 4 = 0$

3.  $x^2 - 2x + 4 = 0$

4.  $x^2 - 4x = 0$

5.  $5r^2 - 3r + 2 = 0$

6.  $7x^2 - 10x - 5 = 0$

7.  $x^2 - 4 = 0$

8.  $25t^2 - 10t = -1$

9.  $6y^2 - 5y = 21$

10.  $2y^2 - 19y = 3$

**Answers: Roots of Quadratic Equations**

1. 1 real root
2. no real roots
3. no real roots
4. 2 real roots
5. no real roots
6. 2 real roots
7. 2 real roots
8. 1 real root
9. 2 real roots
10. 2 real roots