



Comprehensive Summary of Limits and Derivative Calculus

- Limits
- Derivatives
 - Rules, formulas, properties
 - Definition and notation
 - Implicit differentiation
 - Increasing/decreasing
 - Concave up/concave down
 - Extrema
 - Mean value theorem
 - Related rates

Limits

Definitions

Precise Definition : We say $\lim_{x \rightarrow a} f(x) = L$ if for every $\epsilon > 0$ there is a $\delta > 0$ such that whenever $0 < |x - a| < \delta$ then $|f(x) - L| < \epsilon$.

“Working” Definition : We say $\lim_{x \rightarrow a} f(x) = L$ if we can make $f(x)$ as close to L as we want by taking x sufficiently close to a (on either side of a) without letting $x = a$.

Right hand limit : $\lim_{x \rightarrow a^+} f(x) = L$. This has the same definition as the limit except it requires $x > a$.

Left hand limit : $\lim_{x \rightarrow a^-} f(x) = L$. This has the same definition as the limit except it requires $x < a$.

Relationship between the limit and one-sided limits

$$\lim_{x \rightarrow a} f(x) = L \Rightarrow \lim_{x \rightarrow a^+} f(x) = \lim_{x \rightarrow a^-} f(x) = L \quad \lim_{x \rightarrow a} f(x) = L \Rightarrow \lim_{x \rightarrow a} f(x) = L$$

Properties

Assume $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ both exist and c is any number then,

- $\lim_{x \rightarrow a} [cf(x)] = c \lim_{x \rightarrow a} f(x)$
- $\lim_{x \rightarrow a} [f(x) \pm g(x)] = \lim_{x \rightarrow a} f(x) \pm \lim_{x \rightarrow a} g(x)$
- $\lim_{x \rightarrow a} [f(x)g(x)] = \lim_{x \rightarrow a} f(x) \lim_{x \rightarrow a} g(x)$
- $\lim_{x \rightarrow a} \left[\frac{f(x)}{g(x)} \right] = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$ provided $\lim_{x \rightarrow a} g(x) \neq 0$
- $\lim_{x \rightarrow a} [f(x)]^n = \left[\lim_{x \rightarrow a} f(x) \right]^n$
- $\lim_{x \rightarrow a} \sqrt[n]{f(x)} = \sqrt[n]{\lim_{x \rightarrow a} f(x)}$

Basic Limit Evaluations at $\pm \infty$

- Note : $\text{sgn}(a) = 1$ if $a > 0$ and $\text{sgn}(a) = -1$ if $a < 0$.
- $\lim_{x \rightarrow \pm \infty} e^x = \infty$ & $\lim_{x \rightarrow \pm \infty} e^{-x} = 0$
 - $\lim_{x \rightarrow \pm \infty} \ln(x) = \infty$ & $\lim_{x \rightarrow 0^+} \ln(x) = -\infty$
 - If $r > 0$ then $\lim_{x \rightarrow \pm \infty} \frac{b}{x^r} = 0$
 - If $r > 0$ and x^r is real for negative x then $\lim_{x \rightarrow \pm \infty} \frac{b}{x^r} = 0$
 - n even : $\lim_{x \rightarrow \pm \infty} x^n = \infty$
 - n odd : $\lim_{x \rightarrow \pm \infty} x^n = \pm \infty$ & $\lim_{x \rightarrow \pm \infty} x^n = -\infty$
 - n even : $\lim_{x \rightarrow \pm \infty} a x^n + \dots + b x + c = \text{sgn}(a) \infty$
 - n odd : $\lim_{x \rightarrow \pm \infty} a x^n + \dots + b x + c = \text{sgn}(a) \infty$
 - n odd : $\lim_{x \rightarrow \pm \infty} a x^n + \dots + c x + d = -\text{sgn}(a) \infty$

Evaluation Techniques

L'Hospital's Rule

If $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{0}{0}$ or $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\pm \infty}{\pm \infty}$ then,

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

Polynomials at Infinity
 $P(x)$ and $q(x)$ are polynomials. To compute

$$\lim_{x \rightarrow \pm \infty} \frac{P(x)}{q(x)}$$

factor largest power of x out of both $P(x)$ and $q(x)$ and then compute limit.

$$\lim_{x \rightarrow \pm \infty} \frac{3x^2 - 4}{5x - 2x^2} = \lim_{x \rightarrow \pm \infty} \frac{x^2 \left(3 - \frac{4}{x} \right)}{x^2 \left(\frac{5}{x} - 2 \right)} = \lim_{x \rightarrow \pm \infty} \frac{3 - \frac{4}{x}}{\frac{5}{x} - 2} = -\frac{3}{2}$$

Piecewise Function

$$\lim_{x \rightarrow a} g(x) \text{ where } g(x) = \begin{cases} x^2 + 5 & \text{if } x < -2 \\ 1 - 3x & \text{if } x \geq -2 \end{cases}$$

Compute two one sided limits,

$$\lim_{x \rightarrow -2^-} g(x) = \lim_{x \rightarrow -2^-} x^2 + 5 = 9$$

$$\lim_{x \rightarrow -2^+} g(x) = \lim_{x \rightarrow -2^+} 1 - 3x = 7$$

One sided limits are different so $\lim_{x \rightarrow -2} g(x)$ doesn't exist. If the two one sided limits had been equal then $\lim_{x \rightarrow -2} g(x)$ would have existed and had the same value.

Some Continuous Functions

Partial list of continuous functions and the values of x for which they are continuous.

- Polynomials for all x .
- Rational function, except for x 's that give division by zero.
- $\sqrt[n]{x}$ (n odd) for all x .
- $\sqrt[n]{x}$ (n even) for all $x \geq 0$.
- e^x for all x .
- $\ln x$ for $x > 0$.
- $\cos(x)$ and $\sin(x)$ for all x .
- $\tan(x)$ and $\sec(x)$ provided $x \neq \dots, -\frac{3\pi}{2}, -\frac{\pi}{2}, \frac{\pi}{2}, \frac{3\pi}{2}, \dots$
- $\cot(x)$ and $\csc(x)$ provided $x \neq \dots, -2\pi, -\pi, 0, \pi, 2\pi, \dots$

Intermediate Value Theorem

Suppose that $f(x)$ is continuous on $[a, b]$ and let M be any number between $f(a)$ and $f(b)$. Then there exists a number c such that $a < c < b$ and $f(c) = M$.

Derivatives

Definition and Notation

If $y = f(x)$ then the derivative is defined to be $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$.

If $y = f(x)$ then all of the following are equivalent notations for the derivative.

$f'(x) = y' = \frac{df}{dx} = \frac{dy}{dx} = \frac{d}{dx}(f(x)) = Df(x)$ $f'(a) = y'|_{x=a} = \frac{df}{dx}|_{x=a} = \frac{dy}{dx}|_{x=a} = Df(a)$

Interpretation of the Derivative

- If $y = f(x)$ then,
- $m = f'(a)$ is the slope of the tangent line to $y = f(x)$ at $x = a$ and the equation of the tangent line at $x = a$ is given by $y = f(a) + f'(a)(x - a)$.
 - $f'(a)$ is the instantaneous rate of change of $f(x)$ at $x = a$.
 - If $f(x)$ is the position of an object at time x then $f'(a)$ is the velocity of the object at $x = a$.

Basic Properties and Formulas

If $f(x)$ and $g(x)$ are differentiable functions (the derivative exists), c and n are any real numbers,

- $(cf)' = cf'(x)$
- $(f \pm g)' = f'(x) \pm g'(x)$
- $(fg)' = f'g + fg'$ - Product Rule
- $\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$ - Quotient Rule
- $(c)^n = c f'(c) = 0$
- $\frac{d}{dx}(x^n) = nx^{n-1}$ - Power Rule
- $\frac{d}{dx}(f(g(x))) = f'(g(x))g'(x)$ This is the Chain Rule

Common Derivatives

$\frac{d}{dx}(x) = 1$	$\frac{d}{dx}(\csc x) = -\csc x \cot x$	$\frac{d}{dx}(a^x) = a^x \ln(a)$
$\frac{d}{dx}(\sin x) = \cos x$	$\frac{d}{dx}(\cot x) = -\csc^2 x$	$\frac{d}{dx}(e^x) = e^x$
$\frac{d}{dx}(\cos x) = -\sin x$	$\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}$, $x > 0$	$\frac{d}{dx}(\ln(x)) = \frac{1}{x}$, $x > 0$
$\frac{d}{dx}(\tan x) = \sec^2 x$	$\frac{d}{dx}(\cos^{-1} x) = -\frac{1}{\sqrt{1-x^2}}$	$\frac{d}{dx}(\ln x) = \frac{1}{x}$, $x \neq 0$
$\frac{d}{dx}(\sec x) = \sec x \tan x$	$\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$	$\frac{d}{dx}(\log_a(x)) = \frac{1}{x \ln a}$, $x > 0$

Derivatives

Basic Properties/Formulas/Rules

$\frac{d}{dx}(cf(x)) = cf'(x)$, c is any constant. $(f(x) \pm g(x))' = f'(x) \pm g'(x)$
 $\frac{d}{dx}(x^n) = nx^{n-1}$, n is any number. $\frac{d}{dx}(c) = 0$, c is any constant.

$(fg)' = f'g + fg'$ - (Product Rule) $\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$ - (Quotient Rule)

$\frac{d}{dx}(f(g(x))) = f'(g(x))g'(x)$ (Chain Rule)

$\frac{d}{dx}(e^{g(x)}) = g'(x)e^{g(x)}$ $\frac{d}{dx}(\ln g(x)) = \frac{g'(x)}{g(x)}$

Common Derivatives

Polynomials

$\frac{d}{dx}(c) = 0$ $\frac{d}{dx}(x) = 1$ $\frac{d}{dx}(cx) = c$ $\frac{d}{dx}(x^n) = nx^{n-1}$ $\frac{d}{dx}(cx^n) = ncx^{n-1}$

Trig Functions

$\frac{d}{dx}(\sin x) = \cos x$ $\frac{d}{dx}(\cos x) = -\sin x$ $\frac{d}{dx}(\tan x) = \sec^2 x$
 $\frac{d}{dx}(\sec x) = \sec x \tan x$ $\frac{d}{dx}(\csc x) = -\csc x \cot x$ $\frac{d}{dx}(\cot x) = -\csc^2 x$

Inverse Trig Functions

$\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}$ $\frac{d}{dx}(\cos^{-1} x) = -\frac{1}{\sqrt{1-x^2}}$ $\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$
 $\frac{d}{dx}(\sec^{-1} x) = \frac{1}{|x|\sqrt{x^2-1}}$ $\frac{d}{dx}(\csc^{-1} x) = -\frac{1}{|x|\sqrt{x^2-1}}$

Exponential/Logarithm Functions

$\frac{d}{dx}(a^x) = a^x \ln(a)$ $\frac{d}{dx}(e^x) = e^x$ $\frac{d}{dx}(\log_a(x)) = \frac{1}{x \ln a}$, $x > 0$
 $\frac{d}{dx}(\ln(x)) = \frac{1}{x}$, $x > 0$

Hyperbolic Trig Functions

$\frac{d}{dx}(\sinh x) = \cosh x$ $\frac{d}{dx}(\cosh x) = \sinh x$ $\frac{d}{dx}(\tanh x) = \text{sech}^2 x$
 $\frac{d}{dx}(\text{sech } x) = -\text{sech } x \tanh x$ $\frac{d}{dx}(\text{csch } x) = -\text{csch } x \coth x$ $\frac{d}{dx}(\coth x) = -\text{csch}^2 x$

Chain Rule Variants

The chain rule applied to some specific functions.

- $\frac{d}{dx} [f(x)^n] = n[f(x)]^{n-1} f'(x)$
- $\frac{d}{dx} (e^{f(x)}) = f'(x)e^{f(x)}$
- $\frac{d}{dx} (\ln[f(x)]) = \frac{f'(x)}{f(x)}$
- $\frac{d}{dx} (\sin[f(x)]) = f'(x)\cos[f(x)]$
- $\frac{d}{dx} (\cos[f(x)]) = -f'(x)\sin[f(x)]$
- $\frac{d}{dx} (\tan[f(x)]) = f'(x)\sec^2[f(x)]$
- $\frac{d}{dx} (\sec[f(x)]) = f'(x)\sec[f(x)]\tan[f(x)]$
- $\frac{d}{dx} (\tan^{-1}[f(x)]) = \frac{f'(x)}{1+[f(x)]^2}$

Higher Order Derivatives

The n^{th} Derivative is denoted as

$f''(x) = f^{(2)}(x) = \frac{d^2 f}{dx^2}$ and is defined as

$f^{(n)}(x) = (f'(x))'$, i.e. the derivative of first derivative, $f'(x)$.
 $f^{(n)}(x) = (f^{(n-1)}(x))'$, i.e. the derivative of the $(n-1)^{\text{th}}$ derivative, $f^{(n-1)}(x)$.

Implicit Differentiation

Find y' if $e^{2x-9y} + x^3 y^2 + 2x^3 y y' = \cos(y) y' + 11$. Remember $y = y(x)$ here, so products/quotients of x and y will use the product/quotient rule and derivatives of y will use the chain rule. The "trick" is to differentiate as normal and every time you differentiate a y you tack on a y' (from the chain rule). After differentiating solve for y' .

$$e^{2x-9y}(2-9y') + 3x^2 y^2 + 2x^3 y y' = \cos(y) y' + 11$$

$$2e^{2x-9y} - 9y'e^{2x-9y} + 3x^2 y^2 + 2x^3 y y' = \cos(y) y' + 11$$

$$(2x^3 y - 9e^{2x-9y} - \cos(y)) y' = 11 - 2e^{2x-9y} - 3x^2 y^2$$

Critical Points

$x = c$ is a critical point of $f(x)$ provided either

- $f'(c) = 0$ or
- $f'(c)$ doesn't exist.

Increasing/Decreasing

- If $f'(x) > 0$ for all x in an interval I then $f(x)$ is increasing on the interval I .
- If $f'(x) < 0$ for all x in an interval I then $f(x)$ is decreasing on the interval I .
- If $f'(x) = 0$ for all x in an interval I then $f(x)$ is constant on the interval I .

Inflection Points

$x = c$ is an inflection point of $f(x)$ if the concavity changes at $x = c$.

Extrema

Absolute Extrema

- $x = c$ is an absolute maximum of $f(x)$ if $f(c) \geq f(x)$ for all x in the domain.
- $x = c$ is an absolute minimum of $f(x)$ if $f(c) \leq f(x)$ for all x in the domain.

Fermat's Theorem

If $f(x)$ has a relative (or local) extrema at $x = c$, then $x = c$ is a critical point of $f(x)$.

Extreme Value Theorem

If $f(x)$ is continuous on the closed interval $[a, b]$ then there exist numbers c and d so that

- $a \leq c, d \leq b$.
- $f(c)$ is the abs. max. in $[a, b]$.
- $f(d)$ is the abs. min. in $[a, b]$.

Finding Absolute Extrema

To find the absolute extrema of the continuous function $f(x)$ on the interval $[a, b]$ use the following process.

- Find all critical points of $f(x)$ in $[a, b]$.
- Evaluate $f(x)$ at all points found in Step 1.
- Evaluate $f(a)$ and $f(b)$.
- Identify the abs. max. (largest function value) and the abs. min. (smallest function value) from the evaluations in Steps 2 & 3.

Increasing/Decreasing – Concave Up/Concave Down

Concave Up/Concave Down

- If $f''(x) > 0$ for all x in an interval I then $f(x)$ is concave up on the interval I .
- If $f''(x) < 0$ for all x in an interval I then $f(x)$ is concave down on the interval I .

Relative (local) Extrema

- $x = c$ is a relative (or local) maximum of $f(x)$ if $f(c) \geq f(x)$ for all x near c .
- $x = c$ is a relative (or local) minimum of $f(x)$ if $f(c) \leq f(x)$ for all x near c .

1st Derivative Test

If $x = c$ is a critical point of $f(x)$ then $x = c$ is

- a rel. max. of $f(x)$ if $f'(x) > 0$ to the left of $x = c$ and $f'(x) < 0$ to the right of $x = c$.
- a rel. min. of $f(x)$ if $f'(x) < 0$ to the left of $x = c$ and $f'(x) > 0$ to the right of $x = c$.
- not a relative extrema of $f(x)$ if $f'(x)$ is the same sign on both sides of $x = c$.

2nd Derivative Test

If $x = c$ is a critical point of $f(x)$ such that

- $f''(c) = 0$ then $x = c$
- is a relative maximum of $f(x)$ if $f''(c) < 0$.
- is a relative minimum of $f(x)$ if $f''(c) > 0$.
- may be a relative maximum, relative minimum, or neither if $f''(c) = 0$.

Finding Relative Extrema and/or Classifying Critical Points

- Find all critical points of $f(x)$.
- Use the 1st derivative test or the 2nd derivative test on each critical point.

Mean Value Theorem

If $f(x)$ is continuous on the closed interval $[a, b]$ and differentiable on the open interval (a, b) then there is a number $a < c < b$ such that $f'(c) = \frac{f(b) - f(a)}{b - a}$.


Newton's Method

If x_n is the n^{th} guess for the root/solution of $f(x) = 0$ then $(n+1)^{\text{th}}$ guess is $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$ provided $f'(x_n)$ exists.

Related Rates

Sketch picture and identify known/unknown quantities. Write down equation relating quantities and differentiate with respect to t using implicit differentiation (i.e. add on a derivative every time you differentiate a function of t). Plug in known quantities and solve for the unknown quantity.

Ex. A 15 foot ladder is resting against a wall. The bottom is initially 10 ft away and is being pushed towards the wall at $\frac{1}{4}$ ft/sec. How fast is the top moving after 12 sec?




x' is negative because x is decreasing. Using Pythagorean Theorem and differentiating.

$$x^2 + y^2 = 15^2 \Rightarrow 2xx' + 2yy' = 0$$

After 12 sec we have $x = 10 - 12(\frac{1}{4}) = 7$ and so $y = \sqrt{15^2 - 7^2} = \sqrt{176}$. Plug in and solve for y' .

$$7(-\frac{1}{4}) + \sqrt{176} y' = 0 \Rightarrow y' = \frac{7}{4\sqrt{176}} \text{ ft/sec}$$

Ex. Two people are 50 ft apart when one starts walking north. The angle θ changes at 0.01 rad/min. At what rate is the distance between them changing when $\theta = 0.5$ rad?



We have $\theta' = 0.01$ rad/min. and want to find x' . We can use various trig funcs but easiest is,

$$\sec \theta = \frac{x}{50} \Rightarrow \sec \theta \tan \theta \theta' = \frac{x'}{50}$$

We know $\theta = 0.05$ so plug in θ' and solve.


$$\sec(0.5) \tan(0.5)(0.01) = \frac{x'}{50}$$

$x' = 0.3112$ ft/sec
Remember to have calculator in radians!

Optimization

Sketch picture if needed, write down equation to be optimized and constraint. Solve constraint for one of the two variables and plug into first equation. Find critical points of equation in range of variables and verify that they are min/max as needed.

Ex. We're enclosing a rectangular field with 500 ft of fence material and one side of the field is a building. Determine dimensions that will maximize the enclosed area.



Maximize $A = xy$ subject to constraint of $x + 2y = 500$. Solve constraint for x and plug into area.

$$x = 500 - 2y \Rightarrow A = y(500 - 2y) = 500y - 2y^2$$

Differentiate and find critical point(s).


$$A' = 500 - 4y \Rightarrow y = 125$$

By 2nd deriv. test this is a rel. max. and so is the answer we're after. Finally, find x .

$$x = 500 - 2(125) = 250$$

The dimensions are then 250 x 125.

Ex. Determine point(s) on $y = x^2 + 1$ that are closest to $(0, 2)$.



Minimize $f = d^2 = (x-0)^2 + (y-2)^2$ and the constraint is $y = x^2 + 1$. Solve constraint for x^2 and plug into the function.

$$x^2 = y - 1 \Rightarrow f = x^2 + (y - 2)^2$$

Differentiate and find critical point(s).

$$f' = 2y - 3 \Rightarrow y = \frac{3}{2}$$

By the 2nd derivative test this is a rel. min. and so all we need to do is find x value(s).

$$x^2 = \frac{3}{2} - 1 = \frac{1}{2} \Rightarrow x = \pm \frac{1}{\sqrt{2}}$$

The 2 points are then $(\frac{1}{\sqrt{2}}, \frac{3}{2})$ and $(-\frac{1}{\sqrt{2}}, \frac{3}{2})$.