



Comprehensive Summary of Integral Calculus

Integrals

- Rules, Formulas, Properties
- Fundamental Theorem of Calculus
- Integration Techniques
- Area, Volumes of Revolution
- Work
- Arc Length

Integrals
Definitions

Definite Integral: Suppose $f(x)$ is continuous on $[a, b]$. Divide $[a, b]$ into n subintervals of width Δx and choose x_i^* from each interval. Then $\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x$.

Fundamental Theorem of Calculus

Variants of Part I:

Part I: If $f(x)$ is continuous on $[a, b]$ then $g(x) = \int_a^x f(t) dt$ is also continuous on $[a, b]$ and $g'(x) = \frac{d}{dx} \int_a^x f(t) dt = f(x)$.

Part II: $f(x)$ is continuous on $[a, b]$, $F(x)$ is an anti-derivative of $f(x)$ (i.e. $F'(x) = f(x)$) then $\int_a^b f(x) dx = F(b) - F(a)$.

Properties

$\int f(x) \pm g(x) dx = \int f(x) dx \pm \int g(x) dx$
 $\int_a^b f(x) \pm g(x) dx = \int_a^b f(x) dx \pm \int_a^b g(x) dx$
 $\int_a^a f(x) dx = 0$
 $\int_a^b f(x) dx = -\int_b^a f(x) dx$

If $f(x) \geq g(x)$ on $a \leq x \leq b$ then $\int_a^b f(x) dx \geq \int_a^b g(x) dx$
 If $f(x) \geq 0$ on $a \leq x \leq b$ then $\int_a^b f(x) dx \geq 0$
 If $m \leq f(x) \leq M$ on $a \leq x \leq b$ then $m(b-a) \leq \int_a^b f(x) dx \leq M(b-a)$

Common Integrals

$\int k dx = kx + c$
 $\int x^n dx = \frac{1}{n+1} x^{n+1} + c, n \neq -1$
 $\int x^{-1} dx = \int \frac{1}{x} dx = \ln|x| + c$
 $\int \frac{1}{a+x} dx = \frac{1}{a} \ln|ax+b| + c$
 $\int \ln u du = u \ln(u) - u + c$
 $\int e^x dx = e^x + c$
 $\int \tan u du = \ln|\sec u| + c$
 $\int \sec u du = \ln|\sec u + \tan u| + c$
 $\int \frac{1}{a^2+u^2} du = \frac{1}{a} \tan^{-1}(\frac{u}{a}) + c$
 $\int \frac{1}{\sqrt{a^2-u^2}} du = \sin^{-1}(\frac{u}{a}) + c$
 $\int \csc u du = -\cot u + c$
 $\int \sec^2 u du = \tan u + c$

Standard Integration Techniques

Note that at many schools all but the Substitution Rule tend to be taught in a Calculus II class.

u Substitution: The substitution $u = g(x)$ will convert $\int_a^b f(g(x))g'(x) dx = \int_{g(a)}^{g(b)} f(u) du$ using $du = g'(x) dx$. For indefinite integrals drop the limits of integration.

Ex. $\int_1^2 5x^2 \cos(x^3) dx = \int_{\frac{1}{27}}^{\frac{8}{27}} \frac{5}{3} \cos(u) du$
 $u = x^3 \Rightarrow du = 3x^2 dx \Rightarrow x^2 dx = \frac{1}{3} du$
 $x=1 \Rightarrow u=1^3=1 \therefore x=2 \Rightarrow u=2^3=8$
 $= \frac{5}{3} \sin(u) \Big|_{\frac{1}{27}}^{\frac{8}{27}} = \frac{5}{3} (\sin(8) - \sin(1))$

Integration by Parts: $\int u dv = uv - \int v du$ and $\int_a^b u dv = uv \Big|_a^b - \int_a^b v du$. Choose u and dv from integral and compute du by differentiating u and compute v using $v = \int dv$.

Ex. $\int_0^5 x e^{-x} dx$
 $u = x \quad dv = e^{-x} \Rightarrow du = dx \quad v = -e^{-x}$
 $\int x e^{-x} dx = -x e^{-x} + \int e^{-x} dx = -x e^{-x} - e^{-x} + c$

Ex. $\int_3^5 \ln x dx$
 $u = \ln x \quad dv = dx \Rightarrow du = \frac{1}{x} dx \quad v = x$
 $\int_3^5 \ln x dx = x \ln x \Big|_3^5 - \int_3^5 1 dx = (x \ln(x) - x) \Big|_3^5$
 $= 5 \ln(5) - 3 \ln(3) - 2$

Products and (some) Quotients of Trig Functions

For $\int \sin^n x \cos^n x dx$ we have the following:

- n odd. Strip 1 sine out and convert rest to cosines using $\sin^2 x = 1 - \cos^2 x$, then use the substitution $u = \cos x$.
- m odd. Strip 1 cosine out and convert rest to sines using $\cos^2 x = 1 - \sin^2 x$, then use the substitution $u = \sin x$.
- n and m both odd. Use either 1. or 2.
- n and m both even. Use double angle and/or half angle formulas to reduce the integral into a form that can be integrated.

Trig Formulas: $\sin(2x) = 2 \sin(x) \cos(x)$, $\cos^2(x) = \frac{1}{2}(1 + \cos(2x))$, $\sin^2(x) = \frac{1}{2}(1 - \cos(2x))$

Ex. $\int \tan^3 x \sec^2 x dx$
 $\int \tan^3 x \sec^2 x dx = \int \tan^2 x \sec^2 x \tan x \sec x dx$
 $= \int (\sec^2 x - 1) \sec^2 x \tan x \sec x dx$
 $= \int (u^2 - 1) u^2 du \quad (u = \sec x)$
 $= \frac{1}{3} \sec^3 x - \frac{1}{2} \sec^2 x + c$

Ex. $\int \frac{\sin^2 x}{\cos^3 x} dx$
 $\int \frac{\sin^2 x}{\cos^3 x} dx = \int \frac{\sin x \sin x}{\cos^3 x} dx = \int \frac{\sin^2 x \sin x}{\cos^3 x} dx$
 $= \int \frac{(1 - \cos^2 x) \sin x}{\cos^3 x} dx \quad (u = \cos x)$
 $= -\int \frac{(1 - u^2) u^2}{u^3} du = -\int \frac{1 - 2u^2 + u^4}{u^3} du$
 $= \frac{1}{2} \sec^2 x + 2 \ln|\cos x| - \frac{1}{2} \cos^2 x + c$

Inverse Trig Functions

$$\int \frac{1}{\sqrt{a^2 - u^2}} du = \sin^{-1} \left(\frac{u}{a} \right) + c$$

$$\int \frac{1}{a^2 + u^2} du = \frac{1}{a} \tan^{-1} \left(\frac{u}{a} \right) + c$$

$$\int \frac{1}{u\sqrt{u^2 - a^2}} du = \frac{1}{a} \sec^{-1} \left(\frac{u}{a} \right) + c$$

Hyperbolic Trig Functions

$$\int \sinh u \, du = \cosh u + c$$

$$\int \operatorname{sech} u \, du = -\operatorname{csch} u + c$$

$$\int \tanh u \, du = \ln | \cosh u | + c$$

$$\int \operatorname{sech}^2 u \, du = \sinh u + c$$

$$\int \operatorname{csch} u \, du = -\operatorname{csch} u + c$$

$$\int \operatorname{csch}^2 u \, du = -\operatorname{coth} u + c$$

$$\int \operatorname{sech} u \, du = \tan^{-1} | \sinh u | + c$$

Miscellaneous

$$\int \frac{1}{a^2 - u^2} du = \frac{1}{2a} \ln \left| \frac{u+a}{u-a} \right| + c$$

$$\int \sqrt{a^2 + u^2} \, du = \frac{u}{2} \sqrt{a^2 + u^2} + \frac{a^2}{2} \ln | u + \sqrt{a^2 + u^2} | + c$$

$$\int \sqrt{u^2 - a^2} \, du = \frac{u}{2} \sqrt{u^2 - a^2} - \frac{a^2}{2} \ln | u + \sqrt{u^2 - a^2} | + c$$

$$\int \sqrt{a^2 - u^2} \, du = \frac{u}{2} \sqrt{a^2 - u^2} + \frac{a^2}{2} \sin^{-1} \left(\frac{u}{a} \right) + c$$

$$\int \sqrt{2au - u^2} \, du = \frac{u-a}{2} \sqrt{2au - u^2} + \frac{a^2}{2} \cos^{-1} \left(\frac{a-u}{a} \right) + c$$

Standard Integration Techniques

Note that all but the first one of these tend to be taught in a Calculus II class.

u Substitution

Given $\int_a^b f(g(x))g'(x)dx$ then the substitution $u = g(x)$ will convert this into the integral, $\int_a^b f(u)du$.

Integration by Parts

The standard formulas for integration by parts are,

$$\int u \, dv = uv - \int v \, du$$

Choose u and dv and then compute du by differentiating u and compute v by using the fact that $v = \int dv$.

Trig Substitutions

If the integral contains the following root use the given substitution and formula.

$$\sqrt{a^2 - b^2 x^2} \Rightarrow x = \frac{a}{b} \sin \theta \quad \text{and} \quad \cos^2 \theta = 1 - \sin^2 \theta$$

$$\sqrt{b^2 x^2 - a^2} \Rightarrow x = \frac{a}{b} \sec \theta \quad \text{and} \quad \tan^2 \theta = \sec^2 \theta - 1$$

$$\sqrt{a^2 + b^2 x^2} \Rightarrow x = \frac{a}{b} \tan \theta \quad \text{and} \quad \sec^2 \theta = 1 + \tan^2 \theta$$

Partial Fractions

If integrating $\int \frac{P(x)}{Q(x)} dx$ where the degree (largest exponent) of $P(x)$ is smaller than the degree of $Q(x)$ then factor the denominator as completely as possible and find the partial fraction decomposition of the rational expression. Integrate the partial fraction decomposition (P.F.D.). For each factor in the denominator we get term(s) in the decomposition according to the following table.

Factor in $Q(x)$	Term in P.F.D	Factor in $Q(x)$	Term in P.F.D
$ax + b$	$\frac{A}{ax + b}$	$(ax + b)^k$	$\frac{A_1}{ax + b} + \frac{A_2}{(ax + b)^2} + \dots + \frac{A_k}{(ax + b)^k}$
$ax^2 + bx + c$	$\frac{Ax + B}{ax^2 + bx + c}$	$(ax^2 + bx + c)^k$	$\frac{A_1 x + B_1}{ax^2 + bx + c} + \dots + \frac{A_k x + B_k}{(ax^2 + bx + c)^k}$

Products and (some) Quotients of Trig Functions

$$\int \sin^n x \cos^m x \, dx$$

- If n is odd. Strip one sine out and convert the remaining sines to cosines using $\sin^2 x = 1 - \cos^2 x$, then use the substitution $u = \cos x$
- If m is odd. Strip one cosine out and convert the remaining cosines to sines using $\cos^2 x = 1 - \sin^2 x$, then use the substitution $u = \sin x$
- If n and m are both odd. Use either 1. or 2.
- If n and m are both even. Use double angle formula for sine and/or half angle formulas to reduce the integral into a form that can be integrated.

$$\int \tan^n x \sec^m x \, dx$$

- If n is odd. Strip one tangent and one secant out and convert the remaining tangents to secants using $\tan^2 x = \sec^2 x - 1$, then use the substitution $u = \sec x$
- If m is even. Strip two secants out and convert the remaining secants to tangents using $\sec^2 x = 1 + \tan^2 x$, then use the substitution $u = \tan x$
- If n is odd and m is even. Use either 1. or 2.
- If n is even and m is odd. Each integral will be dealt with differently.

Convert Example : $\cos^6 x = (\cos^2 x)^3 = (1 - \sin^2 x)^3$


Trig Substitutions : If the integral contains the following root use the given substitution and formula to convert into an integral involving trig functions.

$$\sqrt{a^2 - b^2 x^2} \Rightarrow x = \frac{a}{b} \sin \theta \quad \sqrt{b^2 x^2 - a^2} \Rightarrow x = \frac{a}{b} \sec \theta \quad \sqrt{a^2 + b^2 x^2} \Rightarrow x = \frac{a}{b} \tan \theta$$

$$\cos^2 \theta = 1 - \sin^2 \theta \quad \tan^2 \theta = \sec^2 \theta - 1 \quad \sec^2 \theta = 1 + \tan^2 \theta$$

Ex. $\int \frac{16}{x^2 \sqrt{4-9x^2}} dx$
 $x = \frac{2}{3} \sin \theta \Rightarrow dx = \frac{2}{3} \cos \theta d\theta$
 $\sqrt{4-9x^2} = \sqrt{4-4\sin^2 \theta} = \sqrt{4\cos^2 \theta} = 2|\cos \theta|$
 Recall $\sqrt{x^2} = |x|$. Because we have an indefinite integral we'll assume positive and drop absolute value bars. If we had a definite integral we'd need to compute θ 's and remove absolute value bars based on that and,
 $|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$
 In this case we have $\sqrt{4-9x^2} = 2 \cos \theta$.

$\int \frac{16}{9 \sin^2 \theta (2 \cos \theta)} d\theta = \int \frac{16}{9 \sin^2 \theta} d\theta = \int \frac{16}{9} \csc^2 \theta d\theta = -\frac{16}{9} \cot \theta + C$
 Use Right Triangle Trig to go back to x 's. From substitution we have $\sin \theta = \frac{3x}{2}$, so,



From this we see that $\cot \theta = \frac{\sqrt{4-9x^2}}{3x}$. So,
 $\int \frac{16}{x^2 \sqrt{4-9x^2}} dx = -\frac{16 \sqrt{4-9x^2}}{3x} + C$

Partial Fractions : If integrating $\int \frac{f(x)}{Q(x)} dx$ where the degree of $f(x)$ is smaller than the degree of $Q(x)$. Factor denominator as completely as possible and find the partial fraction decomposition of the rational expression. Integrate the partial fraction decomposition (P.F.D.). For each factor in the denominator we get term(s) in the decomposition according to the following table.

Factor in $Q(x)$	Term in P.F.D.	Factor in $Q(x)$	Term in P.F.D.
$ax + b$	$\frac{A}{ax + b}$	$(ax + b)^k$	$\frac{A_1}{ax + b} + \frac{A_2}{(ax + b)^2} + \dots + \frac{A_k}{(ax + b)^k}$
$ax^2 + bx + c$	$\frac{Ax + B}{ax^2 + bx + c}$	$(ax^2 + bx + c)^k$	$\frac{A_1 x + B_1}{ax^2 + bx + c} + \dots + \frac{A_k x + B_k}{(ax^2 + bx + c)^k}$

Ex. $\int \frac{7x^2 + 13x}{(x-1)(x^2+4)} dx = \int \frac{A}{x-1} + \frac{Bx+C}{x^2+4} dx$
 $= \int \frac{A}{x-1} + \frac{Bx}{x^2+4} + \frac{C}{x^2+4} dx$
 $= 4 \ln|x-1| + \frac{1}{2} \ln|x^2+4| + 8 \tan^{-1}(\frac{x}{2})$
 Here is partial fraction form and recombined.

$\frac{7x^2+13x}{(x-1)(x^2+4)} = \frac{A}{x-1} + \frac{Bx+C}{x^2+4} = \frac{A(x^2+4) + (Bx+C)(x-1)}{(x-1)(x^2+4)}$
 Set numerators equal and collect like terms.
 $7x^2 + 13x = (A+B)x^2 + (C-B)x + 4A - C$
 Set coefficients equal to get a system and solve to get constants.
 $A+B=7 \quad C-B=13 \quad 4A-C=0$
 $A=4 \quad B=3 \quad C=16$

An alternate method that *sometimes* works to find constants. Start with setting numerators equal in previous example : $7x^2 + 13x = A(x^2 + 4) + (Bx + C)(x - 1)$. Choose nice values of x and plug in. For example if $x = 1$ we get $20 = 5A$ which gives $A = 4$. This won't always work easily.

Applications of Integrals

Net Area : $\int_a^b f(x) dx$ represents the net area between $f(x)$ and the x -axis with area above x -axis positive and area below x -axis negative.



Area Between Curves : The general formulas for the two main cases for each are,

$$y = f(x) \Rightarrow A = \int_a^b [\text{upper function}] - [\text{lower function}] dx \quad \& \quad x = f(y) \Rightarrow A = \int_c^d [\text{right function}] - [\text{left function}] dy$$

If the curves intersect then the area of each portion must be found individually. Here are some sketches of a couple possible situations and formulas for a couple of possible cases.

Volumes of Revolution : The two main formulas are $V = \int A(x) dx$ and $V = \int A(y) dy$. Here is some general information about each method of computing and some examples.

Cylinders

$$A = 2\pi (\text{radius}) (\text{width / height})$$

Rings
 $A = \pi ((\text{outer radius})^2 - (\text{inner radius})^2)$
 Limits: xy of right/bot ring to xy of left/top ring
 Horz. Axis use $f(x)$, Vert. Axis use $f(y)$,
 $g(x), A(x)$ and dx . $g(y), A(y)$ and dy . $g(x), A(x)$ and dx .

Ex. Axis : $y = a > 0$ Ex. Axis : $y = a \leq 0$ Ex. Axis : $y = a > 0$ Ex. Axis : $y = a \leq 0$

outer radius : $a - f(x)$ inner radius : $|a| + f(x)$ radius : $|a| + y$ width : $f(y) - g(y)$

These are only a few cases for horizontal axis of rotation. If axis of rotation is the x -axis use the $y = a \leq 0$ case with $a = 0$. For vertical axis of rotation ($x = a > 0$ and $x = a \leq 0$) interchange x and y to get appropriate formulas.

Work : If a force of $F(x)$ moves an object Average Function Value : The average value of $f(x)$ on $a \leq x \leq b$ is $f_{avg} = \frac{1}{b-a} \int_a^b f(x) dx$
 in $a \leq x \leq b$, the work done is $W = \int_a^b F(x) dx$

Arc Length Surface Area : Note that this is often a Calc II topic. The three basic formulas are,
 $L = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$ if $y = f(x)$, $a \leq x \leq b$ $SA = \int_a^b 2\pi r ds$ (rotate about x-axis) $SA = \int_a^b 2\pi r ds$ (rotate about y-axis)
 where ds is dependent upon the form of the function being worked with as follows.

$ds = \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$ if $y = f(x)$, $a \leq x \leq b$ $ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$ if $x = f(t)$, $y = g(t)$, $a \leq t \leq b$
 $ds = \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$ if $x = f(y)$, $a \leq y \leq b$ $ds = \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$ if $r = f(\theta)$, $a \leq \theta \leq b$

With surface area you may have to substitute in for the x or y depending on your choice of ds to match the differential in the ds . With parametric and polar you will always need to substitute.

Improper Integral

An improper integral is an integral with one or more infinite limits and/or discontinuous integrands. Integral is called convergent if the limit exists and has a finite value and divergent if the limit doesn't exist or has infinite value. This is typically a Calc II topic.

Infinite Limit

- $\int_a^{\infty} f(x) dx = \lim_{t \rightarrow \infty} \int_a^t f(x) dx$ 2. $\int_{-\infty}^b f(x) dx = \lim_{t \rightarrow -\infty} \int_t^b f(x) dx$
- $\int_{-\infty}^{\infty} f(x) dx = \int_{-\infty}^c f(x) dx + \int_c^{\infty} f(x) dx$ provided BOTH integrals are convergent.

Discontinuous Integrals

- Discont. at a : $\int_a^b f(x) dx = \lim_{t \rightarrow a^+} \int_t^b f(x) dx$ 2. Discont. at b : $\int_a^b f(x) dx = \lim_{t \rightarrow b^-} \int_a^t f(x) dx$
- Discontinuity at $a < c < b$: $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$ provided both are convergent.

Comparison Test for Improper Integrals : If $f(x) \geq g(x) \geq 0$ on $[a, \infty)$ then,

- If $\int_a^{\infty} f(x) dx$ conv, then $\int_a^{\infty} g(x) dx$ conv. 2. If $\int_a^{\infty} g(x) dx$ divg, then $\int_a^{\infty} f(x) dx$ divg.

Useful fact : If $a > 0$ then $\int_a^{\infty} \frac{1}{x^p} dx$ converges if $p > 1$ and diverges for $p \leq 1$.

Approximating Definite Integrals

For given integral $\int_a^b f(x) dx$ and a n (must be even for Simpson's Rule) define $\Delta x = \frac{b-a}{n}$ and divide $[a, b]$ into n subintervals $[x_0, x_1], [x_1, x_2], \dots, [x_{n-1}, x_n]$ with $x_0 = a$ and $x_n = b$ then,

Midpoint Rule : $\int_a^b f(x) dx \approx \Delta x [f(x_1^*) + f(x_2^*) + \dots + f(x_n^*)]$, x_i^* is midpoint $[x_{i-1}, x_i]$

Trapezoid Rule : $\int_a^b f(x) dx \approx \frac{\Delta x}{2} [f(x_0) + 2f(x_1) + 2f(x_2) + \dots + 2f(x_{n-1}) + f(x_n)]$

Simpson's Rule : $\int_a^b f(x) dx \approx \frac{\Delta x}{3} [f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + \dots + 4f(x_{n-1}) + f(x_n)]$